



Contribution ID: 34

Type: Poster

Quench of the electronic order in VTe₂ by enhanced lattice fluctuations

Charge-density-wave (CDW) materials, in which electrons and phonons cooperatively interact to form a new symmetry-broken state, stand as ideal candidates to study the mechanisms governing the melting of a macroscopically ordered phase. Furthermore, given their sensitivity to external stimuli, CDW materials constitute a promising platform to investigate the possibility of controlling their properties. Particularly interesting, although little explored, is the case in which the CDW phase transition is determined by the presence of a strong momentum dependent electron-phonon coupling, since it can lead to significant modifications of the properties of these compounds.

TR-ARPES experiments performed on the TMDC compound VTe₂ have revealed that the closure of the CDW gap is not driven by the amplitude modes (AMs) of the system, but is instead dominated by an incoherent process. By applying a so-called three-temperature model (3TM), we demonstrated that the quench of the CDW gap arises as a consequence of the excitation of a subset of strongly-coupled phonon modes which determine a loss of the long-range CDW order. These results therefore suggest that the photoinduced phase transition is determined by non-CDW phonons that interact with the CDW order, thus highlighting the role played by the phonon-phonon interactions.

Primary author: TUNIZ, Manuel (Università degli studi di Trieste)

Co-authors: Dr PUNTEL, Denny (Università degli studi di Trieste); CILENTO, Federico (Elettra - Sincrotrone Trieste SCpA); SAMMARTINO, Francesco (Università degli Studi di Trieste); Prof. PARMIGIANI, Fulvio (Università degli studi di Trieste); BRONSCH, Wibke (Elettra Sincrotrone Trieste)

Presenter: TUNIZ, Manuel (Università degli studi di Trieste)

Session Classification: Session 3